

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 0 688 869 B1**

(12)

## EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention  
of the grant of the patent:  
19.03.2003 Bulletin 2003/12

(51) Int Cl.7: **C12N 15/12, C07K 14/51,  
C12N 5/10, C12N 1/21,  
A61K 38/18**

(21) Application number: **95111771.2**

(22) Date of filing: **30.06.1987**

(54) **Novel osteoinductive compositions**

Osteoinduktive Zusammensetzungen

Compositions ostéoinductrices

**BEST AVAILABLE COPY**

(84) Designated Contracting States:  
**AT BE CH DE FR GB IT LI LU NL SE**

(30) Priority: **01.07.1986 US 880776  
17.12.1986 US 943332  
20.03.1987 US 28285  
26.03.1987 US 31346**

(43) Date of publication of application:  
**27.12.1995 Bulletin 1995/52**

(60) Divisional application:  
**02014841.7 / 1 254 956**

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:  
**87905023.5 / 0 313 578**

(73) Proprietor: **Genetics Institute, LLC**  
**Cambridge, MA 02140 (US)**

(72) Inventors:  
• **Wang, Elizabeth A.**  
**Carlisle, MA 01741 (US)**  
• **Wozney, John M.**  
**Hudson, MA 01749 (US)**  
• **Rosen, Vicki A.**  
**Boston, MA 02116 (US)**

(74) Representative: **VOSSIUS & PARTNER**  
**Postfach 86 07 67**  
**81634 München (DE)**

(56) References cited:  
**US-A- 4 455 256**

**EP 0 688 869 B1**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

## Description

[0001] The present invention relates to novel proteins and processes for obtaining them. These proteins are capable of inducing cartilage and bone formation.

### Background

[0002] Bone is a highly specialized tissue characterized by an extensive matrix structure formed of fibrous bundles of the protein collagen, and proteoglycans, noncollagenous proteins, lipids and acidic proteins. The processes of bone formation and renewal/repair of bone tissue, which occur continuously throughout life, are performed by specialized cells. Normal embryonic long bone development is preceded by formation of a cartilage model. Bone growth is presumably mediated by "osteoblasts" (bone-forming cells), while remodeling of bone is apparently accomplished by the joint activities of bone-resorbing cells, called "osteoclasts" and osteoblasts. A variety of osteogenic, cartilage-inducing and bone inducing factors have been described. See, e.g. European patent applications 148,155 and 169,016 for discussions thereof.

### Brief Description of the Invention

[0003] The present invention provides a novel protein in purified form BMP-3, wherein BMP is bone morphogenic protein. This protein is characterized by peptide sequences the same as or substantially homologous to amino acid sequences illustrated in Tables IV A+B below. It is capable of inducing bone formation at a predetermined site. This bone inductive factor is further characterized by biochemical and biological characteristics including activity at a concentration of 10 to 1000ng/gram of bone in an *in vivo* rat bone formation assay described below. Proteins of this invention may be encoded by the DNA sequences depicted in the Tables or by sequences capable of hybridizing thereto under stringent conditions and coding for polypeptides with bone growth factor biological properties or other variously modified sequences demonstrating such properties.

[0004] The bone inductive factor of the invention, BMP-3, is represented by the bovine homolog bBMP-3. bBMP-3 is characterized by the DNA sequence and amino acid sequence of Table IV A and B which represents the bovine genomic sequence. It is characterized by at least a portion of a peptide sequence the same or substantially the same as amino acid #1 through amino acid #175 of Table IV A and B. BMP-3 is further characterized by the ability to induce bone formation. The bovine factor may be employed as a tool for obtaining the analogous human BMP-3 protein or other mammalian bone inductive proteins. The proper characterization of this bovine bone inductive factor provides the essential "starting point" for the method employing this sequence. The method, employing techniques known to those skilled in the art of genetic engineering, involves using the bovine DNA sequence as a probe to screen a human genomic or cDNA library; and identifying the DNA sequences which hybridize to the probes. A clone with a hybridizable sequence is plaque purified and the DNA isolated therefrom, subcloned and subjected to DNA sequence analysis. Thus another aspect of this invention is a human protein hBMP-3, produced by this method.

[0005] Another aspect of the invention provides pharmaceutical compositions containing a therapeutically effective amount of the bone growth factor polypeptide according to the invention in a pharmaceutically acceptable vehicle. These compositions may further include other therapeutically useful agents. They may also include an appropriate matrix for delivering the proteins to the site of the bone defect and for providing a structure for bone growth. These compositions may be employed in methods for treating a number of bone defects and periodontal disease. These methods, according to the invention, entail administering to a patient needing such bone formation an effective amount of the novel protein BMP-3 as described herein.

[0006] Still a further aspect of the invention are DNA sequences coding on expression for a human or bovine polypeptide having the ability to induce bone formation. Such sequences include the sequence or nucleotides in a 5' to 3' direction illustrated in Tables IV A+B. Alternatively, a DNA sequence which hybridizes under stringent conditions with the DNA sequences of Tables IV A+B and which codes on expression for a protein having at least one bone growth factor biological property are included in the present invention. Finally, allelic or other variations of the sequences of Tables IV A+B, whether such nucleotide changes result in changes in the peptide sequence or not, are also included in the present invention.

[0007] Still a further aspect of the invention is a vector containing a DNA sequence as described above in operative association with an expression control sequence. Such vector may be employed in a novel process for producing a bone growth factor polypeptide in which a cell line transformed with a DNA sequence encoding expression of a bone growth factor polypeptide in operative association with an expression control sequence therefor, is cultured. This claimed process may employ a number of known cells as host cells for expression of the polypeptide. Presently preferred cell lines are mammalian cell lines and bacterial cells.

[0008] Other aspects and advantages of the present invention will be apparent upon consideration of the following

detailed description and preferred embodiments thereof.

### Detailed Description of the Invention

5 [0009] The protein of the present invention is characterized by amino acid sequences or portions thereof the same as or substantially homologous to the sequences shown in Tables IV A+B below. This protein is also characterized by the ability to induce bone formation.

[0010] The bone growth factors provided herein also include factors encoded by the sequences similar to those of Tables IV A+B, but into which modifications are naturally provided (e.g. allelic variations in the nucleotide sequence  
10 which may result in amino acid changes in the polypeptide) or deliberately engineered. For example, synthetic polypeptides may wholly or partially duplicate continuous sequences of the amino acid residues of Tables IV A+B. These sequences, by virtue of sharing primary, secondary, or tertiary structural and conformational characteristics with the bone growth factor polypeptide of Tables IV A+B may possess bone growth factor biological properties in common therewith. Thus, they may be employed as biologically active substitutes for naturally-occurring bone growth factor  
15 polypeptides in therapeutic processes.

[0011] Other specific mutations of the sequences of the bone growth factor described herein involve modifications of one or both of the glycosylation sites. The absence of glycosylation or only partial glycosylation results from amino acid substitution or deletion at one or both of the asparagine-linked glycosylation recognition sites present in the sequences of the bone growth factor shown in Tables IV A+B. The asparagine-linked glycosylation recognition sites  
20 comprise tripeptide sequences which are specifically recognized by appropriate cellular glycosylation enzymes. These tripeptide sequences are either asparagine-X-threonine or asparagine-X-serine, where X is usually any amino acid. A variety of amino acid substitutions or deletions at one or both of the first or third amino acid positions of a glycosylation recognition site (and/or amino acid deletion at the second position) results in non-glycosylation at the modified tripeptide sequence.

[0012] The present invention also encompasses the novel DNA sequences, free of association with DNA sequences encoding other proteinaceous materials, and coding on expression for bone growth factors. These DNA sequences include those depicted in Tables IV A+B in a 5' to 3' direction and those sequences which hybridize under stringent hybridization conditions [see, T. Maniatis et al, Molecular Cloning (A Laboratory Manual), Cold Spring Harbor Laboratory (1982), pages 387 to 389] to the DNA sequences of Tables IV A+B.

[0013] Similarly, DNA sequences which code for bone growth factor polypeptides coded for by the sequences of Tables IV A+B, but which differ in codon sequence due to the degeneracies of the genetic code or allelic variations (naturally-occurring base changes in the species population which may or may not result in an amino acid change) also encode the novel growth factors described herein. Variations in the DNA sequences of Tables IV A+B which are caused by point mutations or by induced modifications to enhance the activity, half-life or production of the polypeptides  
35 encoded thereby are also encompassed in the invention.

[0014] Another aspect of the present invention provides a novel method for producing the novel osteoinductive factors. The method of the present invention involves culturing a suitable cell or cell line, which has been transformed with a DNA sequence coding on expression for a novel bone growth factor polypeptide of the invention, under the control of known regulatory sequences. Suitable cells or cell lines may be mammalian cells, such as Chinese hamster ovary cells (CHO). The selection of suitable mammalian host cells and methods for transformation, culture, amplification, screening and product production and purification are known in the art. See, e.g., Gething and Sambrook, Nature, 293:620-625 (1981), or alternatively, Kaufman et al, Mol. Cell. Biol., 5(7):1750-1759 (1985) or Howley et al, U.S. Patent 4,419,446. Another suitable mammalian cell line, which is described in the accompanying examples, is the monkey COS-1 cell line. A similarly useful mammalian cell line is the CV-1 cell line.

[0015] Bacterial cells are suitable hosts. For example, the various strains of *E. coli* (e.g., HB101, MC1061) are well-known as host cells in the field of biotechnology. Various strains of *B. subtilis*, *Pseudomonas*, other bacilli and the like may also be employed in this method.

[0016] Many strains of yeast cells known to those skilled in the art are also available as host cells for expression of the polypeptides of the present invention. Additionally, where desired, insect cells may be utilized as host cells in the method of the present invention. See, e.g. Miller et al, Genetic Engineering, 8:277-298 (Plenum Press 1986) and references, cited therein.

[0017] Another aspect of the present invention provides vectors for use in the method of expression of these novel osteoinductive polypeptides. Preferably the vectors contain the full novel DNA sequences described above which code for the novel factors of the invention. Additionally the vectors also contain appropriate expression control sequences  
55 permitting expression of the bone inductive protein sequences. Alternatively, vectors incorporating modified sequences as described above are also embodiments of the present invention and useful in the production of the bone inductive proteins. The vectors may be employed in the method of transforming cell lines and contain selected regulatory sequences in operative association with the DNA coding sequences of the invention which are capable of directing the



replication and expression thereof in selected host cells. Useful regulatory sequences for such vectors are known to one of skill in the art and may be selected depending upon the selected host cells. Such selection is routine and does not form part of the present invention.

**[0018]** A protein of the present invention, which induces bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures. An osteogenic preparation employing this protein of the invention may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. De novo bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery. An osteogenic factor of the invention may be valuable in the treatment of periodontal disease, and in other tooth repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of progenitors of bone-forming cells. Of course, the proteins of the invention may have other therapeutic uses.

**[0019]** A further aspect of the invention is a therapeutic method and composition for repairing fractures and other conditions related to bone defects or periodontal diseases. Such a composition comprises a therapeutically effective amount of the bone inductive factor protein of the invention. The bone inductive factor according to the present invention may be present in a therapeutic composition in admixture with a pharmaceutically acceptable vehicle or matrix. Further therapeutic methods and compositions of the invention comprise a therapeutic amount of a bone inductive factor of the invention with a therapeutic amount of at least one of the other bone inductive factors of the invention. Additionally, the protein according to the present invention may be co-administered with one or more different osteoinductive factors with which it may interact. Further, the bone inductive protein may be combined with other agents beneficial to the treatment of the bone defect in question. Such agents include, but are not limited to various growth factors. The preparation of such physiologically acceptable protein compositions, having due regard to pH, isotonicity, stability and the like, is within the skill of the art.

**[0020]** The therapeutic method includes locally administering the composition as an implant or device. When administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone damage. Preferably, the bone growth inductive factor composition would include a matrix capable of delivering the bone inductive factor to the site of bone damage, providing a structure for the developing bone and cartilage and optimally capable of being resorbed into the body. Such matrices may be formed of other materials presently in use for other implanted medical applications.

**[0021]** The choice of material is based on, for example, biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. Similarly, the application of the osteoinductive factors will define the appropriate formulation. Potential matrices for the osteoinductive factors may be biodegradable and chemically defined, such as, but not limited to calcium sulfate, tricalciumphosphate, hydroxyapatite, polylactic acid, polyanhydrides; biodegradable and biologically well defined, such as bone or dermal collagen, other pure proteins or extracellular matrix components; nonbiodegradable and chemically defined, such as sintered hydroxyapatite, bioglass, aluminates, or other ceramics; or combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalciumphosphate. The bioceramics might also be altered in composition, such as in calcium-aluminate-phosphate and processing to alter for example, pore size, particle size, particle shape, and biodegradability.

**[0022]** The dosage regimen will be determined by the attending physician considering various factors which modify the action of such a growth factor, e.g. amount of bone weight desired to be formed, the site of bone damage, the condition of the damaged bone, the patient's age, sex, and diet, the severity of any infection, time of administration and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution. The addition of other known growth factors, such as IGF 1 (insulin like growth factor 1), to the final composition, may also effect the dosage. Generally, the dosage regimen should be in the range of approximately 10 to 10<sup>6</sup> nanograms of protein per gram of bone weight desired. Progress can be monitored by periodic assessment of bone growth and/or repair, e.g. x-rays. Such therapeutic compositions are also presently valuable for veterinary applications due to the lack of species specificity in bone inductive factors. Particularly domestic animals and thoroughbred horses in addition to humans are desired patients for such treatment with the bone inductive factors of the present invention.

**[0023]** The following examples illustrate practice of the present invention in recovering and characterizing the bovine proteins and employing them to recover the human proteins, obtaining the human proteins and in expressing the proteins via recombinant techniques.

#### EXAMPLE I

##### Isolation of Bovine Bone Inductive Factor

**[0024]** Ground bovine bone powder (20-120 mesh, Helitrex) is prepared according to the procedures of M. R. Urist

et al., Proc. Natl Acad. Sci USA, 70:3511 (1973) with elimination of some extraction steps as identified below. Ten kgs of the ground powder is demineralized in successive changes of 0.6N HCl at 4°C over a 48 hour period with vigorous stirring. The resulting suspension is extracted for 16 hours at 4°C with 50 liters of 2M CaCl<sub>2</sub> and 10mM ethylenediamine-tetraacetic acid [EDTA], and followed by extraction for 4 hours in 50 liters of 0.5M EDTA. The residue is washed three times with distilled water before its resuspension in 20 liters of 4M guanidine hydrochloride [GuCl], 20mM Tris (pH 7.4), 1mM N-ethylmaleimide, 1mM iodoacetamide, 1mM phenylmethylsulfonyl fluoride as described in Clin. Orthop. Rel. Res., 171: 213 (1982). After 16 to 20 hours the supernatant is removed and replaced with another 10 liters of GuCl buffer. The residue is extracted for another 24 hours.

[0025] The crude GuCl extracts are combined, concentrated approximately 20 times on a Pellicon apparatus with a 10,000 molecular weight cut-off membrane, and then dialyzed in 50mM Tris, 0.1M NaCl, 6M urea (pH7.2), the starting buffer for the first column. After extensive dialysis the protein is loaded on a 4 liter DEAF cellulose column and the unbound fractions are collected.

[0026] The unbound fractions are concentrated and dialyzed against 50mM NaAc, 50mM NaCl (pH 4.6) in 6M urea. The unbound fractions are applied to a carboxymethyl cellulose column. Protein not bound to the column is removed by extensive washing with starting buffer, and the bone inductive factor containing material desorbed from the column by 50mM NaAc, 0.25mM NaCl, 6M urea (pH 4.6). The protein from this step elution is concentrated 20- to 40- fold, then diluted 5 times with 80mM KPO<sub>4</sub>, 6M urea (pH6.0). The pH of the solution is adjusted to 6.0 with 500mM K<sub>2</sub>HPO<sub>4</sub>. The sample is applied to an hydroxylapatite column (LKB) equilibrated in 80mM KPO<sub>4</sub>, 6M urea (pH6.0) and all unbound protein is removed by washing the column with the same buffer. Bone inductive factor activity is eluted with 100mM KPO<sub>4</sub> (pH7.4) and 6M urea.

[0027] The protein is concentrated approximately 10 times, and solid NaCl added to a final concentration of 0.15M. This material is applied to a heparin - Sepharose column equilibrated in 50mM KPO<sub>4</sub>, 150mM NaCl, 6M urea (pH7.4). After extensive washing of the column with starting buffer, a protein with bone inductive factor activity is eluted by 50mM KPO<sub>4</sub>, 700mM NaCl, 6M urea (pH7.4). This fraction is concentrated to a minimum volume, and 0.4ml aliquots are applied to Superose 6 and Superose 12 columns connected in series, equilibrated with 4M GuCl, 20mM Tris (pH7.2) and the columns developed at a flow rate of 0.25ml/min. The protein demonstrating bone inductive factor activity has a relative migration corresponding to approximately 30,000 dalton protein.

[0028] The above fractions are pooled, dialyzed against 50mM NaAc, 6M urea (pH4.6), and applied to a Pharmacia MonoS HR column. The column is developed with a gradient to 1.0M NaCl, 50mM NaAc, 6M urea (pH4.6). Active fractions are pooled and brought to pH3.0 with 10% trifluoroacetic acid (TFA). The material is applied to a 0.46 x 25cm Vydac C4 column in 0.1% TFA and the column developed with a gradient to 90% acetonitrile, 0.1% TFA (31.5% acetonitrile, 0.1% TFA to 49.5% acetonitrile, 0.1% TFA in 60 minutes at 1ml per minute). Active material is eluted at approximately 40-44% acetonitrile. Aliquots of the appropriate fractions are iodinated by one of the following methods: P. J. McConahey et al, Int. Arch. Allergy, 29:185-189 (1966); A. E. Bolton et al, Biochem J., 133:529 (1973); and D. F. Bowen-Pope, J. Biol. Chem., 237:5161 (1982). The iodinated proteins present in these fractions are analyzed by SDS gel electrophoresis and urea Triton X 100 isoelectric focusing. At this stage, the bone inductive factor is estimated to be approximately 10-50% pure.

## EXAMPLE II

### Characterization of Bovine Bone Inductive Factor

#### A. Molecular Weight

[0029] Approximately 20ug protein from Example I is lyophilized and redissolved in 1X SDS sample buffer. After 15 minutes of heating at 37°C, the sample is applied to a 15% SDS polyacrylamide gel and then electrophoresed with cooling. The molecular weight is determined relative to prestained molecular weight standards (Bethesda Research Labs). Immediately after completion, the gel lane containing bone inductive factor is sliced into 0.3cm pieces. Each piece is mashed and 1.4ml of 0.1% SDS is added. The samples are shaken gently overnight at room temperature to elute the protein. Each gel slice is desalted to prevent interference in the biological assay. The supernatant from each sample is acidified to pH 3.0 with 10% TFA, filtered through a 0.45 micron membrane and loaded on a 0.46cm x 5cm C4 Vydac column developed with a gradient of 0.1% TFA to 0.1% TFA, 90% CH<sub>3</sub>CN. The appropriate bone inductive factor - containing fractions are pooled and reconstituted with 20mg rat matrix. In this gel system, the majority of bone inductive factor fractions have the mobility of a protein having a molecular weight of approximately 28,000 - 30,000 daltons.

## B. Isoelectric Focusing

[0030] The isoelectric point of bone inductive factor activity is determined in a denaturing isoelectric focusing system. The Triton X100 urea gel system (Hoeffer Scientific) is modified as follows: 1) 40% of the ampholytes used are Servalyte 3/10; 60% are Servalyte 7-9. 2) The catholyte used is 40mM NaOH. Approximately 20ug of protein from Example I is lyophilized, dissolved in sample buffer and applied to the isoelectrofocusing gel. The gel is run at 20 watts, 10°C for approximately 3 hours. At completion the lane containing bone inductive factor is sliced into 0.5 cm slices. Each piece is mashed in 1.0ml 6M urea, 5mM Tris (pH 7.8) and the samples agitated at room temperature. The samples are acidified, filtered, desalted and assayed as described above. The major portion of activity as determined in the assay described in Example III migrates in a manner consistent with a pI of 8.8 - 9.2.

## C. Subunit Characterization

[0031] The subunit composition of bone inductive factor is also determined. Pure bone inductive factor is isolated from a preparative 15% SDS gel as described above. A portion of the sample is then reduced with 5mM DTT in sample buffer and re-electrophoresed on a 15% SDS gel. The approximately 30kd protein yields two major bands at approximately 20kd and 18kd, as well as a minor band at 30kd. The broadness of the two bands indicates heterogeneity caused most probably by glycosylation, other post translational modification, proteolytic degradation or carbamylation.

## EXAMPLE III

Biological Activity of Bone Inductive Factor

[0032] A rat bone formation assay according to the general procedure of Sampath and Reddi, Proc. Natl. Acad. Sci. U.S.A., 80:6591-6595 (1983) is used to evaluate the osteogenic activity of the bovine bone inductive factor of the present invention obtained in Example I. This assay can also be used to evaluate bone inductive factors of other species. The ethanol precipitation step is replaced by dialyzing the fraction to be assayed against water. The solution or suspension is then redissolved in a volatile solvent, e.g. 0.1 - 0.2 % TFA, and the resulting solution added to 20mg of rat matrix. This material is frozen and lyophilized and the resulting powder enclosed in #5 gelatin capsules. The capsules are implanted subcutaneously in the abdominal thoracic area of 21 - 49 day old male long Evans rats. The implants are removed after 7-14 days. Half of each implant is used for alkaline phosphatase analysis [See, A. H. Reddi et al., Proc. Natl Acad Sci., 69:1601 (1972)] and half is fixed and processed for histological analysis. Routinely, lum glycolmethacrylate sections are stained with Von Kossa and acid fuchsin to detect new bone mineral. Alkaline phosphatase, an enzyme produced by chondroblasts and osteoblasts in the process of matrix formation, is also measured. New cartilage and bone formation often correlates with alkaline phosphatase levels. Table I below illustrates the dose response of the rat matrix samples including a control not treated with bone inductive factor.

TABLE 1

Protein* Implanted ug	Cartilage	Alk. Phos.u/l
7.5	2	Not done
2.5	3	445.7
0.83	3	77.4
0.28	0	32.5
0.00	0	31.0

\*At this stage the bone inductive factor is approximately 10-15% pure.

[0033] The bone or cartilage formed is physically confined to the space occupied by the matrix. Samples are also analyzed by SDS gel electrophoresis and isoelectric focusing as described above, followed by autoradiography. Analysis reveals a correlation of activity with protein bands at 28 - 30kd and a pI 9.0. An extinction coefficient of 1 OD/mg-cm is used as an estimate for protein and approximating the purity of bone inductive factor in a particular fraction. In the in vivo rat bone formation assays on dilutions as described above, the protein is active in vivo at 10 to 200ng protein/gram bone to probably greater than 1ug protein/gram bone.



EXAMPLE IVBovine Bone Inductive Factor Protein Composition

5 [0034] The protein composition of Example IIA of molecular weight 28 - 30kd is reduced as described in Example IIC and digested with trypsin. Eight tryptic fragments are isolated by standard procedures having the following amino acid sequences:

Fragment 1: A A F L G D I A L D E E D L G  
 10 Fragment 2: A F Q V Q Q A A D L  
 Fragment 3: N Y Q D M V V E G  
 Fragment 4: S T P A Q D V S R  
 Fragment 5: N Q E A L R  
 Fragment 6: L S E P D P S H T L E E  
 15 Fragment 7: F D A Y Y  
 Fragment 8: L K P S N ? A T I Q S I V E

[0035] A less highly purified preparation of protein from bovine bone is prepared according to a purification scheme similar to that described in Example I. The purification basically varies from that previously described by omission of  
 20 the DE-52 column, the CM cellulose column and the mono S column, as well as a reversal in the order of the hydroxylapatite and heparin sepharose columns. Briefly, the concentrated crude 4 M extract is brought to 85% final concentration of ethanol at 4 degrees. The mixture is then centrifuged, and the precipitate redissolved in 50 mM Tris, 0.15 M NaCl, 6.0 M urea. This material is then fractionated on Heparin Sepharose as described. The Heparin bound material is fractionated on hydroxyapatite as described. The active fractions are pooled, concentrated, and fractionated on a  
 25 high resolution gel filtration (TSK 30000 in 6 M guanidinium chloride, 50 mM Tris, pH 7.2). The active fractions are pooled, dialyzed against 0.1% TFA, and then fractionated on a C4 Vydac reverse phase column as described. The preparation is reduced and electrophoresed on an acrylamide gel. The protein corresponding to the 18K band is eluted and digested with trypsin. Tryptic fragments are isolated having the following amino acid sequences:

30 Fragment 9: S L K P S N H A T I Q S ? V  
 Fragment 10: S F D A Y Y C S ? A  
 Fragment 11: V Y P N M T V E S C A  
 Fragment 12: V D F A D I ? W

35 [0036] Tryptic Fragments 7 and 8 are noted to be substantially the same as Fragments 10 and 9, respectively.

A. bBMP-3

40 [0037] Probes consisting of pools of oligonucleotides are designed on the basis of the amino acid sequences of the tryptic Fragments 9 (Probe #3), 10 (Probe #2), and 11 (Probe #1), and synthesized on an automated DNA synthesizer.

Probe #1: A C N G T C A T [A/G] T T N G G [A/G] T A

45 Probe #2: C A [A/G] T A [A/G] T A N G C [A/G] T C [A/G] A A

Probe #3: T G [A/G/T] A T N G T N G C [A/G] T G [A/G] T T

50 [0038] A bovine genomic recombinant library is constructed as follows: Bovine liver DNA is partially digested with the restriction endonuclease enzyme Sau 3A and sedimented through a sucrose gradient. Size fractionated DNA in the range of 15-30kb is then ligated to the bacteriophage Bam HI vector EMBL3 [Frischauf et al, *J. Mol. Biol.*, 170: 827-842 (1983)]. The library is plated at 8000 recombinants per plate. Duplicate nitrocellulose replicas of the plaques  
 55 are made and amplified according to a modification of the procedure of Woo et al, *Proc. Natl. Acad. Sci. USA*, 75: 3688-91 (1978).

[0039] This recombinant bovine genomic library constructed in EMBL3 is screened by the TMAC hybridization procedure, i.e. kyloridized in 3M tetramethylammonium chloride (TMAC), 0.1M sodium phosphate pH6.5, 1mM EDTA, 5X

Denhardt's, 0.6% SDS, 100ug/ml salmon sperm DNA at 48 degrees C, and washed in 3M TMAC, 50mM Tris pH8.0 at 50 degrees C. These conditions minimize the detection of mismatches to the 17 mer probe pool [see, Wood et al, *Proc. Natl. Acad. Sci. U.S.A.*, 82:1585-1588 (1985)]. 400,000 recombinants are screened in duplicate with Probe #1 which has been labeled with <sup>32</sup>P. All recombinants which hybridized to this probe are replated for secondaries. Triplicate nitrocellulose replicas are made of the secondary plates, and amplified as described. The three sets of filters are hybridized to Probes #1, #2 and #3, again under TMAC conditions. One clone, lambda bP-819, hybridizes to all three probes and is plaque purified and DNA is isolated from a plate lysate. Bacteriophage lambda bP-819 was deposited with the ATCC on June 16, 1987 under accession number 40344. This bP-819 clone encodes the bovine bone growth factor designated bBMP-3.

[0040] The region of bP-819 which hybridizes to Probe #2 is localized and sequenced. The partial DNA and derived amino acid sequences of this region are shown in Table IVA. The amino acid sequences corresponding to tryptic Fragments 10 and 12 are underlined. The first underlined sequence corresponds to Fragment 12 while the second corresponds to Fragment 10. This region of bP-819, therefore, which hybridizes to Probe #2 encodes at least 111 amino acids. This amino acid sequence is encoded by the DNA sequence from nucleotide #414 through #746.

TABLE IV. A.

383	393	403	413	(1)	428
GAGGAGGAAG	OGGTCTAAGG	GGGTCTTCT	GOCTCTGCAG	AAC AAT GAG CTT CCT GGG GCA	
				Asn Asn Glu Leu Pro Gly Ala	
443	458	473	488		
GAA TAT CAG TAC AAG GAG GAT GAA GTA TGG GAG GAG AGG AAG CCT TAC AAG ACT					
Glu Tyr Gln Tyr Lys Glu Asp Glu Val Trp Glu Glu Arg Lys Pro Tyr Lys Thr					
503	518	533			
CTT CAG ACT CAG CCC CCT GAT AAG AGT AAG AAC AAA AAG AAA CAG AGG AAG GGA					
Leu Gln Thr Gln Pro Pro Asp Lys Ser Lys Asn Lys Lys Lys Gln Arg Lys Gly					
548	563	578	593		
CCT CAG CAG AAG AGT CAG ACG CTC CAG TTT GAT GAA CAG ACC CTG AAG AAG GCA					
Pro Gln Gln Lys Ser Gln Thr Leu Gln Phe Asp Glu Gln Thr Leu Lys Lys Ala					
608	623	638			
AGA AGA AAG CAA TGG ATT GAA CCC OGG AAT TGT GCC AGA OGG TAC CTT AAA GTG					
Arg Arg Lys Gln Trp Ile Glu Pro Arg Asn Cys Ala Arg Arg Tyr Leu Lys <u>Val</u>					
653	668	683	698		
GAC TTC GCA GAT ATT GGC TGG AGC GAA TGG ATT ATT TOC OCC AAG TOC TTC GAT					
<u>Asp Phe Ala Asp Ile Gly Trp</u> Ser Glu Trp Ile Ile Ser Pro Lys <u>Ser Phe Asp</u>					
713	728	743 (111)	756		
GCC TAT TAC TGC TOC GGA GCG TGC CAG TTC CCC ATG CCA AAG GTAGOCATTG					
<u>Ala Tyr Tyr Cys Ser Gly Ala</u> Cys Gln Phe Pro MET Pro Lys					
766	776	786			
TTTTTTGTGC	TGTCTTCCC	ATTTCATAG			

[0041] The region of bP-819 which hybridizes to Probe #1 and #3 is localized and sequenced. The partial DNA and derived amino acid sequences of this region are shown in Table IVB. The amino acid sequences corresponding to tryptic Fragments 9 and 11 are underlined. The first underlined sequence corresponds to Fragment 9 while the second underlined sequence corresponds to Fragment 11. The peptide sequence of this region of bP-819 which hybridizes to Probe #1 and #3 is 64 amino acids in length encoded by nucleotide #305 through #493 of Table IVB. The arginine



residue encoded by the AGA triplet is presumed to be the carboxy-terminus of the protein based on the presence of a stop codon (TAA) adjacent to it. The nucleic acid sequence preceding the couplet TC (positions 305-306) is presumed to be an intron (non-coding sequence) based on the presence of a consensus acceptor sequence (i.e. a pyrimidine-rich stretch, TTCTCCCTTTTCGTTCT, followed by AG) and the presence of a stop rather than a basic residue in the appropriate position of the derived amino acid sequence.

[0042] bBMP-3 is therefore characterized by the DNA and amino acid sequence of Table IV A and Table IV B. The peptide sequence of this clone is 175 amino acids in length and is encoded by the DNA sequence from nucleotide #414 through nucleotide #746 of Table IV A and nucleotide #305 through nucleotide #493 of Table IV B.

TABLE IV. B.

284	294	304	(112)	319
CTAAOCTGTG	TTCTOOC TTT	TOGTTOCTAG	TCT TTG AAG OCA TCA AAT CAC GCT ACC	Ser Leu Lys Pro Ser Asn His Ala Thr
334	349	364	379	
ATC CAG AGT ATA GTG AGA GCT GTG GGG GTC GTC OCT GGA ATC CCC GAG CCT TGC				
Ile Gln Ser Ile Val	Arg Ala Val Gly Val Val	Pro Gly Ile Pro	Glu Pro Cys	
394	409	424	439	
TGT GTG CCA GAA AAG ATG TOC TCA CTC AGC ATC TTA TTC TTT GAT GAA AAC AAG				
Cys Val Pro Glu Lys MET Ser Ser Leu Ser Ile Leu Phe Phe Asp Glu Asn Lys				
454	469	484	(175)	
AAT GTG GTA CTT AAA GTA TAT CCA AAC ATG ACA GTA GAG TCT TGT GCT TGC AGA				
Asn Val Val Leu Lys Val Tyr Pro Asn MET Thr Val Glu Ser Cys Ala Cys Arg				
503	513	523	533	
TAAOCTGGTG	AAGAACTCAT	CTGGATGCCT	AACTCAATCG	

## EXAMPLE V

## Human BMP-3

[0043] The sequences of BMP-3 as shown in Tables IV A+B have significant homology to the beta (B) and beta (A) subunits of the inhibins. The inhibins are a family of hormones which are presently being investigated for use in contraception. See, A. J. Mason et al, Nature, 318:659-663 (1985). To a lesser extent they are also homologous to Mullerian inhibiting substance (MIS), a testicular glycoprotein that causes regression of the Mullerian duct during development of the male embryo and transforming growth factor-beta (TGF-b) which can inhibit or stimulate growth of cells or cause them to differentiate.

[0044] Because bovine and human bone growth factor genes are presumed to be significantly homologous, oligonucleotide probes which have been shown to hybridize to the bovine DNA sequence of Table IV.A and IV.B are used to screen a human genomic library. A human genomic library (Toole et al.; supra) is screened using these probes, and presumptive positives are isolated and DNA sequence obtained as described above. Evidence that this recombinant encodes a portion of the human bone inductive factor molecule relies on the bovine/human protein and gene structure homologies.

[0045] Once a recombinant bacteriophage containing DNA encoding a portion of the human BMP-3 molecule is obtained the human coding sequence is used as a probe as described in Example V (A) to identify a human cell line or tissue which synthesizes BMP-3. mRNA is selected by oligo (dT) cellulose chromatography and cDNA is synthesized and cloned in lambda gt10 by established techniques (Toole et al., supra).

[0046] Alternatively, the entire gene encoding this human bone inductive factor can be identified and obtained in additional recombinant clones if necessary. Additional recombinants containing further 3' or 5' regions of this human

bone inductive factor gene can be obtained by identifying unique DNA sequences at the end(s) of the original clone and using these as probes to rescreen the human genomic library. The gene can then be reassembled in a single plasmid by standard molecular biology techniques and amplified in bacteria. The entire human BMP-3 factor gene can then be transferred to an appropriate expression vector. The expression vector containing the gene is then transfected into a mammalian cell, e.g. monkey COS cells, where the human gene is transcribed and the RNA correctly spliced. Media from the transfected cells are assayed for bone inductive factor activity as described herein as an indication that the gene is complete. mRNA is obtained from these cells and cDNA synthesized from this mRNA source and cloned. The procedures described above may similarly be employed to isolate other species' bone inductive factor of interest by utilizing the bovine bone inductive factor and/or human bone inductive factor as a probe source. Such other species' bone inductive factor may find similar utility in, inter alia, fracture repair.

## EXAMPLE VI

### Expression of Bone Inductive Factors.

[0047] In order to produce bovine, human or other mammalian bone inductive factors, the DNA encoding it is transferred into an appropriate expression vector and introduced into mammalian cells by conventional genetic engineering techniques.

[0048] One skilled in the art can construct mammalian expression vectors by employing the sequence of Tables IV A+B or other modified sequences and known vectors, such as pCD [Okayama et al., Mol. Cell Biol., 2:161-170 (1982)] and pJL3, pJL4 [Gough et al., EMBO J., 4:645-653 (1985)]. The transformation of these vectors into appropriate host cells can result in expression of osteoinductive factors. One skilled in the art could manipulate the sequences of Tables IV A+B by eliminating or replacing the mammalian regulatory sequences flanking the coding sequence with bacterial sequences to create bacterial vectors for intracellular or extracellular expression by bacterial cells. For example, the coding sequences could be further manipulated (e.g., ligated to other known linkers or modified by deleting non-coding sequences therefrom or altering nucleotides therein by other known techniques). The modified bone inductive factor coding sequence could then be inserted into a known bacterial vector using procedures such as described in T. Taniguchi et al., Proc. Natl Acad. Sci. USA, 77:5230-5233 (1980). This exemplary bacterial vector could then be transformed into bacterial host cells and bone inductive factor expressed thereby. For a strategy for producing extracellular expression of bone inductive factor in bacterial cells, see, e.g. European patent application EPA 177,343.

[0049] Similar manipulations can be performed for the construction of an insect vector [See, e.g. procedures described in published European patent application 155,476] for expression in insect cells. A yeast vector could also be constructed employing yeast regulatory sequences for intracellular or extracellular expression of the factors of the present invention by yeast cells. [See, e.g., procedures described in published PCT application WO86/00639 and European patent application EPA 123,289].

[0050] A method for producing high levels of an osteoinductive factor of the invention from mammalian cells involves the construction of cells containing multiple copies of the heterologous bone inductive factor gene. The heterologous gene can be linked to an amplifiable marker, e.g. the dihydrofolate reductase (DHFR) gene for which cells containing increased gene copies can be selected for propagation in increasing concentrations of methotrexate (MTX) according to the procedures of Kaufman and Sharp, J. Mol. Biol., 159:601-629 (1982). This approach can be employed with a number of different cell types.

[0051] For example, a plasmid containing a DNA sequence for a bone inductive factor of the invention in operative association with other plasmid sequences enabling expression thereof and the DHFR expression plasmid pAdA26SV (A)3 [Kaufman and Sharp, Mol. Cell. Biol., 2:1304 (1982)] can be co-introduced into DHFR-deficient CHO cells, DUKX-BII, by calcium phosphate coprecipitation and transfection. DHFR expressing transformants are selected for growth in alpha media with dialyzed fetal calf serum, and subsequently selected for amplification by growth in increasing concentrations of MTX (sequential steps in 0.02, 0.2, 1.0 and 5uM MTX) as described in Kaufman et al., Mol Cell Biol., 5:1750 (1983). Transformants are cloned, and biologically active bone inductive factor expression is monitored by rat bone formation assay. Bone inductive factor expression should increase with increasing levels of MTX resistance. Similar procedures can be followed to produce other bone inductive factors.

[0052] Alternatively, the human gene is expressed directly, as described above. Active bone inductive factor may be produced in bacteria or yeast cells. However the presently preferred expression system for biologically active recombinant human bone inductive factor is stably transformed CHO cells.

Example VIIBiological Activity of Expressed Bone Inductive Factor

5 [0053] To measure the biological activity of the expressed bone inductive factor obtained in Example VI above. The factor is partially purified on a Heparin Sepharose column. 4 ml of transfection supernatant from one 100 mm dish is concentrated approximately 10 fold by ultrafiltration on a YM 10 membrane and then dialyzed against 20mM Tris, 0.15 M NaCl, pH 7.4 (starting buffer). This material is then applied to a 1.1 ml Heparin Sepharose column in starting buffer. Unbound proteins are removed by an 8 ml wash of starting buffer, and bound proteins are desorbed by a 3-4 ml wash  
10 of 20 mM Tris, 2.0 M NaCl, pH 7.4.

[0054] The proteins bound by the Heparin column are concentrated approximately 10-fold on a Centricon 10 and the salt reduced by diafiltration with 0.1% trifluoroacetic acid. The appropriate amount of this solution is mixed with 20 mg of rat matrix and then assayed for in vivo bone and cartilage formation as previously described in Example III. A mock transfection supernatant fractionation is used as a control.

15 [0055] The implants containing rat matrix to which specific amounts of human BMP-3 have been added are removed from rats after seven days and processed for histological evaluation. Representative sections from each implant are stained for the presence of new bone mineral with von Kossa and acid fuchsin, and for the presence of cartilage-specific matrix formation using toluidine blue. The types of cells present within the section, as well as the extent to which these cells display phenotype are evaluated.

20 [0056] The procedures described above may be employed to isolate other bone inductive factors of interest by utilizing the bovine bone inductive factors and/or human bone inductive factors as a probe source. Such other bone inductive factors may find similar utility in, inter alia, fracture repair.

25 **Claims**

**Claims for the following Contracting States : BE, CH, LI, DE, FR, GB, IT, LU, NL, SE**

30 1. A gene encoding bovine BMP-3 comprising the following DNA sequence:

35

40

45

50

55

383 393 403 413 423  
 GAGGAGGAG CCGTCACCG CCGTCCTCTT CCGTCACCG AAC AAT GAG CTT CTT GGG GCA  
 5 Asn Asn Glu Leu Pro Gly Ala  
  
 443 458 473 488  
 GAA TAT CAG TAC AAG CAG GAT GAA GGA TCG GAG GAG AGG AAG CTT TAC AAG ACT  
 10 Glu Tyr Gln Tyr Lys Glu Asp Glu Val Trp Glu Glu Arg Lys Pro Tyr Lys Thr  
  
 503 513 533  
 CTT CAG ACT CAG CCC CTT GAT AAG AGT AAG AAC AAA AAG AAA CAG AGG AAG GCA  
 Leu Gln Thr Gln Pro Pro Asp Lys Ser Lys Asn Lys Lys Lys Gln Arg Lys Gly  
  
 548 563 578 593  
 CTT CAG CAG AAG AGT CAG ACG CTC CAG TTT GAT GAA CAG ACC CTG AAG AAG GCA  
 15 Pro Gln Gln Lys Ser Gln Thr Leu Gln Phe Asp Glu Gln Thr Leu Lys Lys Ala  
  
 608 623 638  
 AGA AGA AAG CAA TCG ATT GAA CCC CCG AAT TGT GCC AGA CCG TAC CTT AAA GTG  
 20 Arg Arg Lys Gln Trp Ile Glu Pro Arg Asn Cys Ala Arg Arg Tyr Leu Lys Val  
  
 653 668 683 698  
 GAC TTC GCA GAT ATT GCC TCG AGC GAA TCG ATT ATT TCC CCC AAG TCC TTC GAT  
 25 Asp Phe Ala Asp Ile Gly Trp Ser Glu Trp Ile Ile Ser Pro Lys Ser Phe Asp  
  
 713 723 743 756 766  
 GGC TAT TAC TGC TCC GGA GCG TGC CAG TTC CCC AAG CCA AAG CAGACCTTC TTTTTCGCC  
 30 Ala Tyr Tyr Cys Ser Gly Ala Cys Gln Phe Pro Met Pro Lys  
  
 776 786  
 TGTCTTCCCTTCC ATTCCCTAG ; and  
  
 284 294 304 319  
 35 CTAACCTGTC TTCTCCCTTT TCGTCCCTAG TCT TTG AAG CCA TCA AAT CAC GGT ACC  
 Ser Leu Lys Pro Ser Asn His Ala Thr  
  
 334 349 364 379  
 40 ATC CAG AAT ATA GTC AGA GGT GTG GGG GTC GTC CTT GGA ATC CCC GAG CTT TGC  
 Ile Gln Ser Ile Val Arg Ala Val Gly Val Val Pro Gly Ile Pro Glu Pro Cys  
  
 394 409 424 439  
 45 TGT GTG CCA GAA AAG ATG TCC TCA CTC AGC ATC TTA TTC TTT GAT GAA AAC AAG  
 Cys Val Pro Glu Lys Met Ser Ser Leu Ser Ile Leu Phe Phe Asp Glu Asn Lys

2. A gene encoding bovine BMP-3 having the amino acid sequence given in claim 1.

3. A gene encoding a protein exhibiting at least the property of BMP-3 to induces the formation of bone and comprising a DNA sequence:

- (a) which differs from a DNA sequence of claim 1 in codon sequence due to the degeneracy of the genetic code;
- (b) which hybridizes with a DNA sequence of claim 1 or section (a), above, under stringent hybridization conditions; or
- (c) which represents a fragment, or allelic variation of a DNA sequence of claim. 1.



454 459 464  
 AAT GTG GGA CTT AAA GGA TAT CCA AAC ATG ACA GGA GAG TCT TGT GGT TGC AAG  
 5 Asn Val Val Leu Lys Val Tyr Pro Asn Met Thr Val Glu Ser Cys Ala Cys Arg  
 503 513 523 533  
 TACCTGGTG AAGAACTGAT CTGGATGCTT AACTGATCG.

4. The DNA sequence of claim 3, which encodes human BMP-3.
5. The DNA sequence of claim 3 or 4, which is a genomic DNA sequence.
6. The DNA sequence of claim 3 or 4, which is a cDNA sequence.
7. A vector containing the gene or DNA sequence of any one of claims 1 to 6 in operative association with an expression control sequence.
8. A cell transformed with a vector of claim 7.
9. The cell of claim 8 which is a mammalian cell, a bacterial cell, an insect cell or a yeast cell.
10. The cell of claim 9 which is a CHO cell.
11. A protein exhibiting properties of BMP-3 which is encoded by the gene or DNA sequence of any one of claims 1 to 6.
12. A protein exhibiting properties of BMP-3 which is produced by the steps of culturing in a suitable culture medium a cell transformed with an expression vector comprising a gene or a DNA sequence of any one of claims 1 to 6, and recovering said protein from said culture medium.
13. A process for producing the protein of claim 11 or 12, comprising the steps of culturing in a suitable culture medium the cell of claim 9 and isolating said protein from said culture medium.
14. A pharmaceutical composition comprising the protein of claim 11 or 12 and a pharmaceutically acceptable vehicle.
15. The pharmaceutical composition of Claim 14, further comprising a matrix capable of delivering the composition to the site of the bone or cartilage defect and providing a structure for inducing bone or cartilage formation.
16. The pharmaceutical composition of claim 15, wherein said matrix comprises hydroxyapatite, collagen, polylactic acid or tricalcium phosphate.
17. Use of a protein of claim 11 or 12 for the preparation of a pharmaceutical composition for inducing bone or cartilage formation.

#### Claims for the following Contracting State : AT

1. A method for the preparation of a gene encoding bovine BMP-3 comprising the following DNA sequence:

383 393 403 413 423  
 GAGGAGGAG CCGTCACG GGGTCCTTCT GGGTCGAC AAC AAT GAG CTT CTT GGG GCA  
 Asn Asn Glu Leu Pro Gly Ala  
 5  
 443 453 473 483  
 GAA TTT CAG TAC AAG GAG GAT GAA GGA TGG GAG GAG ACG AAG CTT TAC AAG ACT  
 Glu Tyr Gln Tyr Lys Glu Asp Glu Val Trp Glu Glu Arg Lys Pro Tyr Lys Thr  
 10  
 503 513 533  
 CTT CAG ACT CAG CCC CTT GAT AAG AAT AAG AAC AAA AAG AAA CAG ACG AAG GCA  
 Leu Gln Thr Gln Pro Pro Asp Lys Ser Lys Asn Lys Lys Lys Gln Arg Lys Gly  
 15  
 543 563 573 593  
 CTT CAG CAG AAG AAT CAG ACG CTC CAG TTT GAT GAA CAG ACC CTT AAG AAG GCA  
 Pro Gln Gln Lys Ser Gln Thr Leu Gln Phe Asp Glu Gln Thr Leu Lys Lys Ala  
 20  
 603 623 633  
 AGA AGA AAG CAA TGG AAT GAA CCC CCG AAT TGT GGC AGA CCG TAC CTT AAA GTG  
 Arg Arg Lys Gln Trp Ile Glu Pro Arg Asn Cys Ala Arg Arg Tyr Leu Lys Val  
 25  
 653 653 663 693  
 GAC TTC GCA GAT AAT GGC TGG ACC GAA TGG AAT AAT TCC CCC AAG TCC TTC GAT  
 Asp Phe Ala Asp Ile Gly Trp Ser Glu Trp Ile Ile Ser Pro Lys Ser Phe Asp  
 30  
 713 723 743 753 766  
 GGC TAT TAC TGC TCC GGA GCG TGC CAG TTC CCC AAG CCA AAG GAGGCTG TTTTGGCC-  
 Ala Tyr Tyr Cys Ser Gly Ala Cys Gln Phe Pro Met Pro Lys  
 35  
 776 786  
 TGTCTTCCC ATTGCTAG ; and  
 284 294 304 319  
 CTAACCTGCT TTTCTCTTT TGGTTCCTG TCT TTG AAG CCA TCA AAT CAC GGT ACC  
 Ser Leu Lys Pro Ser Asn His Ala Thr  
 334 349 354 379  
 ATC CAG AAT AAT GTG AGA GGT GTG GGG GTG GTG CTT GGA ATC CCC GAG CTT TGC  
 Ile Gln Ser Ile Val Arg Ala Val Gly Val Val Pro Gly Ile Pro Glu Pro Cys  
 394 409 424 439  
 TGT GTG CCA GAA AAG ATG TCC TCA CTC ACC ATC TTA TTC TTT GAT GAA AAC AAG  
 Cys Val Pro Glu Lys Met Ser Ser Leu Ser Ile Leu Phe Phe Asp Glu Asn Lys

2. The method of claim 1, wherein the gene encoding the BMP-3 has the amino acid sequence given in claim 1.
3. The method of claim 1, wherein a gene encoding a protein exhibiting at least the property of BMP-3 to induce the formation of bone comprises a DNA sequence:
  - (a) which differs from a DNA sequence obtained according to the method of claim 1 in codon sequence due to the degeneracy of the genetic code;
  - (b) which hybridises with a DNA sequence obtained according to the method of claim 1 or section (a), above under stringent hybridisation conditions; or
  - (c) which represents a fragment, or allelic variation of a DNA sequence obtained according to the method of claim 1.

454 459 454  
 AAT GAG GGA CTT AAA GGA TAT CCA AAC AIG AGA GGA GAG TCT TGT GGT TGC AGA  
 Asn Val Val Leu Lys Val Tyr Pro Asn Met Thr Val Glu Ser Cys Ala Cys Arg  
 503 513 523 533  
 TACCTGGG AAGAACTCT CTGATGCTT AACGATCG.

said method comprising screening of a recombinant bovine genomic library with probes consisting of pools of oligonucleotides designed on the basis of partial amino acid sequences and isolating a positive clone.

4. The method of claim 3, wherein the DNA sequence encodes human BMP-3.

5. The method of claim 3 or 4, wherein the DNA sequence is a genomic DNA sequence.

6. The method of claim 3 or 4, wherein the DNA sequence is a cDNA sequence.

7. A method for the preparation of a recombinant vector comprising inserting the gene or DNA sequence obtained according to a method of any one of claims 1 to 6 in operative association with an expression control sequence in a suitable vector.

8. A cell transformed with a vector prepared according to the method of claim 7.

9. The cell of claim 8 which is a mammalian cell, a bacterial cell, an insect cell or a yeast cell.

10. The cell of claim 9 which is a CHO cell.

11. A process for producing a protein exhibiting properties of BMP-3 comprising the steps of culturing in a suitable culture medium the cell of claim 9 and isolating said protein from said culture medium.

12. A method for the preparation of a pharmaceutical composition comprising combining the protein produced according to the method of claim 11 with a pharmaceutically acceptable vehicle.

13. The method of claim 12, wherein the pharmaceutical composition further comprises a matrix capable of delivering the composition to the site of the bone or cartilage defect and providing a structure for inducing bone or cartilage formation.

14. The method of claim 13, wherein said matrix of said pharmaceutical composition comprises hydroxyapatite, collagen, polylactic acid or tricalcium phosphate.

15. Use of a protein of claim 11 for the preparation of a pharmaceutical composition for inducing bone or cartilage formation.

16. A gene encoding bovine BMP-3 comprising the following DNA sequence:

5

383 393 403 413 423  
GAGGAGGAG CCGTCTACCG CCGTCTCTCT CCGTCTCGAG AAC AAT GAG CTT CTT GCG GCA  
Asn Asn Glu Leu Pro Gly Ala

10

443 453 473 483  
GAA TTT CAG TAC AAG GAG GAT GAA GAA TGG GAG GAG AGG AAG CTT TAC AAG ACT  
Glu Tyr Gln Tyr Lys Glu Asp Glu Val Trp Glu Glu Arg Lys Pro Tyr Lys Thr

15

503 513 533  
CTT CAG ACT CAG CCC CTT GAT AAG AGT AAG AAC AAA AAG AAA CAG AGG AAG GCA  
Leu Gln Thr Gln Pro Pro Asp Lys Ser Lys Asn Lys Lys Lys Gln Arg Lys Gly

20

548 563 573 593  
CCT CAG CAG AAG AGT CAG ACG CTC CAG TTT GAT GAA CAG ACC CTC AAG AAG GCA  
Pro Gln Gln Lys Ser Gln Thr Leu Gln Phe Asp Glu Gln Thr Leu Lys Lys Ala

25

608 623 638  
AGA AGA AAG CAA TGG ATT GAA CCC CCG AAT TGT GCG AGA CCG TAC CTT AAA GTC  
Arg Arg Lys Gln Trp Ile Glu Pro Arg Asn Cys Ala Arg Arg Tyr Leu Lys Val

30

653 658 683 693  
GAC TTC GCA GAT ATT GGC TGG AGC GAA TGG AAT ATT TCC CCC AAG TCC TTC GAT  
Asp Phe Ala Asp Ile Gly Trp Ser Glu Trp Ile Ile Ser Pro Lys Ser Phe Asp

35

713 723 743 755 765  
GCC TAT TAC TGC TCC GGA GCG TGC CAG TTC CCC AAG CCA AAG GAGGAGTCTG TTTTCTGCTG  
Ala Tyr Tyr Cys Ser Gly Ala Cys Gln Phe Pro Met Pro Lys

40

776 786  
TCTCTTCC CATTCTCTG ; and

45

284 294 304 319  
CTACCTGTC TCTCTCTTT TCTCTCTG TCT TCG AAG CCA TCA AAT CAC GGT ACC  
Ser Leu Lys Pro Ser Asn His Ala Thr

50

334 349 364 379  
ATC CAG AGT AAT GTG AGA GGT GTG GCG GTC GTC CTT GGA ATC CCC GAG CTT TGC  
Ile Gln Ser Ile Val Arg Ala Val Gly Val Val Pro Gly Ile Pro Glu Pro Cys

55

394 409 424 439  
TGT GTG CCA GAA AAG ATG TCC TCA CTC AGC ATC TTA TTC TTT GAT GAA AAC AAG  
Cys Val Pro Glu Lys Met Ser Ser Leu Ser Ile Leu Phe Phe Asp Glu Asn Lys

17. A gene encoding bovine BMP-3 having the amino acid sequence given in claim 16.
18. A gene encoding a protein exhibiting at least the property of BMP-3 to induce the formation of bone and comprising a DNA sequence:
  - (a) which differs from a DNA sequence of claim 16 in codon sequence due to the degeneracy of the genetic code;
  - (b) which hybridises with a DNA sequence of claim 16 or section (a), above, under stringent hybridisation conditions; or
  - (c) which represents a fragment, or allelic variation of a DNA sequence of claim 16.



454 469 484  
 AAT GAG GGA CTT AAA GGA TAT CCA AAC AAG ACG GGA GAG TGT TGT GGT TGC AAA  
 Asn Val Val Leu Lys Val Tyr Pro Asn Met Thr Val Glu Ser Cys Ala Cys Arg  
 5  
 503 513 523 533  
 TACCTGTC AAGACTCT CTGGTCTT AACTGATC.  
 10

19. The DNA sequence of claim 18, which encodes human BMP-3.

20. The DNA sequence of claim 18 or 19, which is a genomic DNA sequence.

21. The DNA sequence of claim 18 or 19, which is a cDNA sequence.

22. A vector containing the gene or DNA sequence of any one of claims 16 to 21 in operative association with an expression control sequence.

23. A cell transformed with a vector of claim 22.

24. The cell of claim 23 which is a mammalian cell, a bacterial cell, an insect cell or a yeast cell.

25. The cell of claim 24, which is a CHO cell.

26. A protein exhibiting properties of BMP-3 which is encoded by the gene or DNA sequence of any one of claims 1 to 21.

27. A protein exhibiting properties of BMP-3 which is produced by the steps of culturing in a suitable culture medium a cell transformed with an expression vector comprising a gene or a DNA sequence of any one of claims 1 to 21 and recovering said protein from said culture medium.

28. A pharmaceutical composition comprising the protein of claim 26 or 27 and a pharmaceutically acceptable vehicle.

29. The pharmaceutical composition of claim 29, further comprising a matrix capable of delivering the composition to the site of the bone or cartilage defect and providing a structure for inducing bone or cartilage formation.

30. The pharmaceutical composition of claim 29, wherein said matrix comprises hydroxyapatite, collagen, polylactic acid or tricalcium phosphate.

## Patentansprüche

Patentansprüche für folgende Vertragsstaaten : BE, CH, LI, DE, FR, GB, IT, LU, NL, SE

1. Gen, das bovines BMP-3 codiert, welches die folgende DNA-Sequenz umfasst:

363 393 403 413 426  
 GAGGAGAG GAGTCTGAG GAGTCTCTCT GAGTCTGAG AAC AAT GAG CTT CTT GGG GGA  
 Asn Asn Glu Leu Pro Gly Ala

5 443 458 473 488  
 GAA TTT CAG TAC AAG GAG GAT GAA GGA TGG GAG GAG AAG AAG CTT TAC AAG ACT  
 Glu Tyr Gln Tyr Lys Glu Asp Glu Val Trp Glu Glu Arg Lys Pro Tyr Lys Thr

10 503 513 533  
 CTT CAG ACT CAG CCC COT GAT AAG AAT AAG AAC AAA AAG AAA CAG AGG AAG GGA  
 Leu Gln Thr Gln Pro Pro Asp Lys Ser Lys Asn Lys Lys Lys Gln Arg Lys Gly

15 548 563 578 593  
 CTT CAG CAG AAG AAT CAG AAG CTC CAG TTT GAT GAA CAG ACC CTG AAG AAG GGA  
 Pro Gln Gln Lys Ser Gln Thr Leu Gln Phe Asp Glu Gln Thr Leu Lys Lys Ala

20 608 623 638  
 AAG AAG AAG CAA TGG AAT GAA CCC CAG AAT TGT GCG AAG CAG CTT AAA GTC  
 Arg Arg Lys Gln Trp Ile Glu Pro Arg Asn Cys Ala Arg Arg Tyr Leu Lys Val

25 653 668 683 693  
 GAC TTC GGA GAT AAT GGC TGG AAG GAA TGG AAT AAT TCC CCC AAG TCC TTC GAT  
 Asp Phe Ala Asp Ile Gly Trp Ser Glu Trp Ile Ile Ser Pro Lys Ser Phe Asp

713 723 743 755 766  
 GGC TAT TAC TGC TCC GGA GCG TGC CAG TTC CCC ATG CCA AAG GAGGCTG TTTTGTCT-  
 Ala Tyr Tyr Cys Ser Gly Ala Cys Gln Phe Pro Met Pro Lys

776 786  
 TGTCTTCCC ATTCTCTG ; and

30 284 294 304 319  
 CTAACCTGTC TTTCTCTTT TCTTCTGAG TCT TTC AAG CCA TCA AAT CAC GGT ACC  
 Ser Leu Lys Pro Ser Asn His Ala Thr

35 334 349 364 379  
 ATC CAG AAT AAT GTC AAG GGT GTC GGC GTC GTC CTT GGA ATC CCC GAG CTT TGC  
 Ile Gln Ser Ile Val Arg Ala Val Gly Val Val Pro Gly Ile Pro Glu Pro Cys

40 394 409 424 439  
 TGT GTC CCA GAA AAG ATG TCC TCA CTC AAG ATC TTA TTC TTT GAT GAA AAC AAG  
 Cys Val Pro Glu Lys Met Ser Ser Leu Ser Ile Leu Phe Phe Asp Glu Asn Lys

45 454 469 484  
 AAT GTC GGA CTT AAA GGA TAT CCA AAC ATG AAG GGA CAG TCT TGT GGT TGC AAG  
 Asn Val Val Leu Lys Val Tyr Pro Asn Met Thr Val Glu Ser Cys Ala Cys Arg

503 513 523 533  
 TACCTGTC AAGACTCT CTGCTGCTT AACTGATC

2. Gen, das bovines BMP-3 codiert, welches die Aminosäuresequenz aufweist, die in Anspruch 1 angegeben ist.
3. Gen, das ein Protein codiert, das mindestens die Eigenschaft von BMP-3 aufweist, die Bildung von Knochen zu induzieren, und eine DNA-Sequenz umfasst:

(a) die sich von einer DNA-Sequenz nach Anspruch 1 in der Codonsequenz auf Grund der Degeneriertheit des genetischen Codes unterscheidet;

- (b) die mit einer DNA-Sequenz nach Anspruch 1 oder Absatz (a) oben unter stringenten Hybridisierungsbedingungen hybridisiert; oder  
 (c) die ein Fragment oder eine allelische Variation einer DNA-Sequenz nach Anspruch 1 darstellt.

- 5 4. DNA-Sequenz nach Anspruch 3, die menschliches BMP-3 codiert.
5. DNA-Sequenz nach Anspruch 3 oder 4, die eine genomische DNA-Sequenz ist.
6. DNA-Sequenz nach Anspruch 3 oder 4, die eine cDNA-Sequenz ist.
- 10 7. Vektor, der das Gen oder die DNA-Sequenz nach einem der Ansprüche 1 bis 6 in funktioneller Verknüpfung mit einer Expressionskontrollsequenz enthält.
8. Zelle, transformiert mit einem Vektor nach Anspruch 7.
- 15 9. Zelle nach Anspruch 8, die eine Säugierzelle, eine Bakterienzelle, eine Insektenzelle oder eine Hefezelle ist.
10. Zelle nach Anspruch 9, die eine CHO-Zelle ist.
- 20 11. Protein, das Eigenschaften von BMP-3 aufweist, das von dem Gen oder der DNA-Sequenz nach einem der Ansprüche 1 bis 6 codiert wird.
12. Protein, das Eigenschaften von BMP-3 aufweist, das durch die Schritte des Züchtens in einem geeigneten Kulturmedium von einer Zelle, die mit einem Expressionsvektor transformiert ist, der ein Gen oder eine DNA-Sequenz nach einem der Ansprüche 1 bis 6 umfasst, und des Gewinnens des Proteins aus dem Kulturmedium hergestellt wird.
- 25 13. Verfahren zur Herstellung des Proteins nach Anspruch 11 oder 12, umfassend die Schritte des Züchtens der Zelle nach Anspruch 9 in einem geeigneten Kulturmedium und des Isolierens des Proteins aus dem Kulturmedium.
- 30 14. Arzneimittel, umfassend das Protein nach Anspruch 11 oder 12 und ein pharmazeutisch verträgliches Vehikel.
15. Arzneimittel nach Anspruch 14, das zusätzlich eine Matrix umfasst, die das Mittel an die Stelle des Knochen- oder Knorpeldefekts bringen und eine Struktur zum Induzieren von Knochen- oder Knorpelbildung zur Verfügung stellen kann.
- 35 16. Arzneimittel nach Anspruch 15, wobei die Matrix Hydroxyapatit, Kollagen, Polymilchsäure oder Tricalciumphosphat umfasst.
- 40 17. Verwendung eines Proteins nach Anspruch 11 oder 12 für die Herstellung eines Arzneimittels zum Induzieren von Knochen- oder Knorpelbildung.

#### Patentansprüche für folgenden Vertragsstaat : AT

- 45 1. Verfahren zur Herstellung eines Gens, das bovines BMP-3 codiert, welches die folgende DNA-Sequenz umfasst:
- 50
- 55

383 393 403 413 425  
 GAGGAGGAG CCGTCACG CCGTCCTCT CCGTCCTCG · AAC AAT GAG CTT CTT GCG GCA  
 Asn Asn Glu Leu Pro Gly Ala  
 5  
 443 453 473 486  
 GAA TAT CAG TAC AAG GAG GAT GAA GGA TCG GAG GAG ACG AAG CTT TAC AAG ACT  
 Glu Tyr Gln Tyr Lys Glu Asp Glu Val Trp Glu Glu Arg Lys Pro Tyr Lys Thr  
 10  
 503 513 533  
 CTT CAG ACT CAG CCC COT GAT AAG AAT AAG AAC AAA AAG AAA CAG ACG AAG GGA  
 Leu Gln Thr Gln Pro Pro Asp Lys Ser Lys Asn Lys Lys Lys Gln Arg Lys Gly  
 15  
 548 563 573 583  
 CTT CAG CAG AAG AAT CAG ACG CTC CAG TTT GAT GAA CAG ACC CTG AAG AAG GCA  
 Pro Gln Gln Lys Ser Gln Thr Leu Gln Phe Asp Glu Gln Thr Leu Lys Lys Ala  
 20  
 608 623 638  
 AGA AGA AAG CAA TCG AAT GAA CCC CCG AAT TGT GCG AGA CCG TAC CTT AAA GTC  
 Arg Arg Lys Gln Trp Ile Glu Pro Arg Asn Cys Ala Arg Arg Tyr Leu Lys Val  
 25  
 653 668 683 693  
 GAC TTC GCA GAT AAT GGC TCG ACC GAA TCG AAT AAT TCC CCC AAG TCC TTC GAT  
 Asp Phe Ala Asp Ile Gly Trp Ser Glu Trp Ile Ile Ser Pro Lys Ser Phe Asp  
 30  
 713 723 743 755 766  
 GGC TAT TAC TGC TCC GGA GCG TCC CAG TTC CCC AAG CCA AAG GAGGAGTTC TTTTTCCTCC--  
 Ala Tyr Tyr Cys Ser Gly Ala Cys Gln Phe Pro Met Pro Lys  
 35  
 776 786  
 TGTCTTCCC ATTTCCTAG ; and  
 284 294 304 319  
 CTAACCTGTC TTTCTCTTT TCGTTCTAG TCT TTC AAG CCA TCA AAT CAC GGT ACC  
 Ser Leu Lys Pro Ser Asn His Ala Thr  
 35  
 334 349 364 379  
 ATC CAG AAT AAT GTC AGA GGT GTC GGG GTC GTC CTT GGA ATC CCC GAG CTT TCC  
 Ile Gln Ser Ile Val Arg Ala Val Gly Val Val Pro Gly Ile Pro Glu Pro Cys  
 40  
 394 409 424 439  
 TGT GTC CCA GAA AAG ATG TCC TCA CTC ACC ATC TTA TTC TTT GAT GAA AAC AAG  
 Cys Val Pro Glu Lys Met Ser Ser Leu Ser Ile Leu Phe Phe Asp Glu Asn Lys  
 45  
 454 469 484  
 AAT GTC GGA CTT AAA GGA TAT CCA AAC ATG ACG GGA GAG TCT TGT GGT TCC AGA  
 Asn Val Val Leu Lys Val Tyr Pro Asn Met Thr Val Glu Ser Cys Ala Cys Arg  
 50  
 503 513 523 533  
 TACCTGTC AAGACTCT CTGATCTT AACTGATC  
 55  
 2. Verfahren nach Anspruch 1, wobei das Gen, das BMP-3 codiert, die Aminosäuresequenz aufweist, die in Anspruch 1 angegeben ist.  
 3. Verfahren nach Anspruch 1, wobei ein Gen, das ein Protein codiert, welches mindestens die Eigenschaft von BMP-



3 zeigt, die Bildung von Knochen zu induzieren, eine DNA-Sequenz umfasst:

- (a) die sich von einer DNA-Sequenz, die gemäß dem Verfahren nach Anspruch 1 erhalten wurde, in der Codonsequenz auf Grund der Degeneriertheit des genetischen Codes unterscheidet;
- (b) die mit einer DNA-Sequenz, die gemäß dem Verfahren nach Anspruch 1 oder Absatz (a) oben erhalten wurde, unter stringenten Hybridisierungsbedingungen hybridisiert; oder
- (c) die ein Fragment oder eine allelische Variation einer DNA-Sequenz darstellt, die gemäß dem Verfahren nach Anspruch 1 erhalten wurde.

4. Verfahren nach Anspruch 3, wobei die DNA-Sequenz menschliches BMP-3 codiert.

5. Verfahren nach Anspruch 3 oder 4, wobei die DNA-Sequenz eine genomische DNA-Sequenz ist.

6. Verfahren nach Anspruch 3 oder 4, wobei die DNA-Sequenz eine cDNA-Sequenz ist.

7. Verfahren zur Herstellung eines rekombinanten Vektors, umfassend das Einfügen des Gens oder der DNA-Sequenz, die gemäß einem Verfahren nach einem der Ansprüche 1 bis 6 erhalten wurde, in funktioneller Verknüpfung mit einer Expressionskontrollsequenz in einen geeigneten Vektor.

8. Zelle, transformiert mit einem Vektor, der gemäß dem Verfahren nach Anspruch 7 hergestellt wurde.

9. Zelle nach Anspruch 8, die eine Säugierzelle, eine Bakterienzelle, eine Insektenzelle oder eine Hefezelle ist.

10. Zelle nach Anspruch 9, die eine CHO-Zelle ist.

11. Verfahren zur Herstellung eines Proteins, das Eigenschaften von BMP-3 aufweist, umfassend die Schritte des Züchtens der Zelle nach Anspruch 9 in einem geeigneten Kulturmedium und des Isolierens des Proteins aus dem Kulturmedium.

12. Verfahren zur Herstellung eines Arzneimittels, umfassend das Kombinieren des Proteins, das gemäß dem Verfahren nach Anspruch 11 hergestellt wurde, mit einem pharmazeutisch verträglichen Vehikel.

13. Verfahren nach Anspruch 12, wobei das Arzneimittel zusätzlich eine Matrix umfasst, die das Mittel an die Stelle des Knochen- oder Knorpeldefekts bringen und eine Struktur für das Induzieren von Knochen- oder Knorpelbildung zur Verfügung stellen kann.

14. Verfahren nach Anspruch 13, wobei die Matrix des Arzneimittels Hydroxyapatit, Kollagen, Polymilchsäure oder Tricalciumphosphat umfasst.

15. Verwendung eines Proteins nach Anspruch 11 für die Herstellung eines Arzneimittels zum Induzieren von Knochen- oder Knorpelbildung.

16. Gen, das bovines BMP-3 codiert, welches die folgende DNA-Sequenz umfasst:

383 393 403 413 423  
 GAGGAGGAG CAGTCCGAG GGTGCTTCT GGTCTGCGA AAC AAT GAG CTT CTT GGG GCA  
 Asn Asn Glu Leu Pro Gly Ala  
 5  
 443 453 473 483  
 GAA TAT CAG TAC AAG GAG GAT GAA GGA TGG CAG GAG AGG AAG CTT TAC AAG ACT  
 Glu Tyr Gln Tyr Lys Glu Asp Glu Val Trp Glu Glu Arg Lys Pro Tyr Lys Thr  
 10  
 503 513 533  
 CTT CAG ACT CAG CCC CTT GAT AAG AAT AAG AAC AAA AAG AAA CAG AGG AAG GCA  
 Leu Gln Thr Gln Pro Pro Asp Lys Ser Lys Asn Lys Lys Lys Gln Arg Lys Gly  
 15  
 543 563 573 593  
 CTT CAG CAG AAG AAT CAG AAG CTC CAG TTT GAT GAA CAG ACC CTG AAG AAG GCA  
 Pro Gln Gln Lys Ser Gln Thr Leu Gln Phe Asp Glu Gln Thr Leu Lys Lys Ala  
 20  
 603 623 633  
 ACA ACA AAG CAA TGG AAT GAA CCC CAG AAT TGT GGC ACA CAG TAC CTT AAA GTG  
 Arg Arg Lys Gln Trp Ile Glu Pro Arg Asn Cys Ala Arg Arg Tyr Leu Lys Val  
 25  
 653 663 683 693  
 GAC TTC GCA GAT AAT GGC TGG AGC GAA TGG AAT AAT TCC CCC AAG TCC TTC GAT  
 Asp Phe Ala Asp Ile Gly Trp Ser Glu Trp Ile Ile Ser Pro Lys Ser Phe Asp  
 30  
 713 723 743 753 763  
 GGC TAT TAC TCC TCC GGA GGC TCC CAG TTC CCC AAG CCA AAG GAGGCTG TTTTGTGCT  
 Ala Tyr Tyr Cys Ser Gly Ala Cys Gln Phe Pro Met Pro Lys  
 35  
 776 786  
 TGTCTTCCC ATTCTCTG ; and  
 284 294 304 319  
 CTAACCTGAG TTTCTCTT TGTCTCTG TCT TTC AAG CCA TCA AAT CAC GGT ACC  
 Ser Leu Lys Pro Ser Asn His Ala Thr  
 36  
 334 349 354 379  
 ATC CAG AAT AAT GTC AAT GGT GTC GGC GTC GTC CTT GGA ATC CCC GAG CTT TGC  
 Ile Gln Ser Ile Val Arg Ala Val Gly Val Val Pro Gly Ile Pro Glu Pro Cys  
 40  
 394 409 424 439  
 TGT GTC CCA GAA AAG ATC TCC TCA CTC AGC ATC TTA TTC TTT GAT GAA AAC AAG  
 Cys Val Pro Glu Lys Met Ser Ser Leu Ser Ile Leu Phe Phe Asp Glu Asn Lys  
 45  
 454 469 484  
 AAT GTC GGA CTT AAA GGA TAT CCA AAC AAG AGC GGA GAG TCT TGT CTT TGC AAT  
 Asn Val Val Leu Lys Val Tyr Pro Asn Met Thr Val Glu Ser Cys Ala Cys Arg  
 50  
 503 513 523 533  
 TACCTGAG AAGACTCT CTGCTCTT TACTCTTC  
 55

17. Gen, das bovines BMP-3 codiert, welches die Aminosäuresequenz aufweist, die in Anspruch 16 angegeben ist.

18. Gen, das ein Protein codiert, das mindestens die Eigenschaft von BMP-3 aufweist, die Bildung von Knochen zu induzieren, und eine DNA-Sequenz umfasst:

- (a) die sich von einer DNA-Sequenz nach Anspruch 16 in der Codonsequenz auf Grund der Degeneriertheit des genetischen Codes unterscheidet;
- (b) die mit einer DNA-Sequenz nach Anspruch 16 oder Absatz (a) oben unter stringenten Hybridisierungsbe-

dingungen hybridisiert; oder

(c) die ein Fragment oder eine allelische Variation einer DNA-Sequenz nach Anspruch 16 darstellt.

19. DNA-Sequenz nach Anspruch 18, die menschliches BMP-3 codiert.

20. DNA-Sequenz nach Anspruch 18 oder 19, die eine genomische DNA-Sequenz ist.

21. DNA-Sequenz nach Anspruch 18 oder 19, die eine cDNA-Sequenz ist.

22. Vektor, der das Gen oder die DNA-Sequenz nach einem der Ansprüche 16 bis 21 in funktioneller Verknüpfung mit einer Expressionskontrollsequenz enthält.

23. Zelle, transformiert mit einem Vektor nach Anspruch 22.

24. Zelle nach Anspruch 23, die eine Säugerzelle, eine Bakterienzelle, eine Insektenzelle oder eine Hefezelle ist.

25. Zelle nach Anspruch 24, die eine CHO-Zelle ist.

26. Protein, das Eigenschaften von BMP-3 aufweist, das von dem Gen oder der DNA-Sequenz nach einem der Ansprüche 1 bis 21 codiert wird.

27. Protein, das Eigenschaften von BMP-3 aufweist, das durch die Schritte des Züchtens in einem geeigneten Kulturmedium von einer Zelle, die mit einem Expressionsvektor transformiert ist, der ein Gen oder eine DNA-Sequenz nach einem der Ansprüche 1 bis 21 umfasst, und des Gewinnens des Proteins aus dem Kulturmedium hergestellt wird.

28. Arzneimittel, das das Protein nach Anspruch 26 oder 27 und ein pharmazeutisch verträgliches Vehikel umfasst.

29. Arzneimittel nach Anspruch 29, das zusätzlich eine Matrix umfasst, die das Mittel an die Stelle des Knochen- oder Knorpeldefekts bringen und eine Struktur zum Induzieren von Knochen- oder Knorpelbildung zur Verfügung stellen kann.

30. Arzneimittel nach Anspruch 29, wobei die Matrix Hydroxyapatit, Kollagen, Polymilchsäure oder Tricalciumphosphat umfasst.

## Revendications

Revendications pour les Etats contractants suivants : BE, CH, LI, DE, FR, GB, IT, LU, NL, SE

1. Gène codant pour la BMP-3 bovine comprenant la séquence d'ADN suivante :

383 393 403 413 423  
 GAGGAGAG GAGGAGAG GAGGAGAG GAGGAGAG AAC AAT GAG CTT CTT GCG GGA  
 Asn Asn Glu Leu Pro Gly Ala  
 5  
 443 453 473 483  
 GAA TGT CAG TAC AAG GAG GAT GAA GGA TGT GAG GAG AAG AAG CTT TAC AAG ACT  
 Glu Tyr Glu Tyr Lys Glu Asp Glu Val Thr Glu Glu Arg Lys Pro Tyr Lys Thr  
 10  
 503 513 533  
 CTT CAG ACT CAG CCG CCG GAT AAG AAT AAG AAG AAA AAG AAA CAG AAG AAG GGA  
 Leu Glu Thr Glu Pro Pro Asp Lys Ser Lys Asn Lys Lys Lys Glu Arg Lys Gly  
 15  
 543 563 573 583  
 CTT CAG CAG AAG AAT CAG AAG CTC CAG TTT GAT GAA CAG AAG CTT AAG AAG GGA  
 Pro Glu Glu Lys Ser Glu Thr Leu Glu Phe Asp Glu Glu Thr Leu Lys Lys Ala  
 20  
 603 623 633  
 AAG AAG AAG CAA TGT AAT GAA CCG CCG AAT TGT GAT AAG CCG TAC CTT AAA GTC  
 Arg Arg Lys Glu Thr Ile Glu Pro Arg Asn Cys Ala Arg Arg Tyr Leu Lys Val  
 25  
 653 663 683 693  
 GAC TTC GGA GAT AAT GCG TGG AAG GAA TGG AAT AAT TCC CCG AAG TCC TTC GAT  
 Asp Phe Ala Asp Ile Gly Thr Ser Glu Thr Ile Ile Ser Pro Lys Ser Phe Asp  
 713 723 743 753 763  
 GGC TAT TAC TGT TCC GGA CCG TCC CAG TTC CCG AAG CCA AAG GAGGAGAG TTTTGGCC  
 Ala Tyr Tyr Cys Ser Gly Ala Cys Glu Phe Pro Met Pro Lys  
 776 783  
 TGTGTTGCTG ATTGCTG ; et  
 30  
 284 294 304 319  
 CTAACCTGAG TTTGCTG TGTGCTG TGT TGT AAG CCA TCA AAT CAC GAT ACC  
 Ser Leu Lys Pro Ser Asn His Ala Thr  
 35  
 334 349 354 373  
 ATC CAG AAT AAT GTC AAG GAT GTG GCG GTC GTC CTT GGA ATC CCG GAG CTT TCC  
 Ile Glu Ser Ile Val Arg Ala Val Gly Val Val Pro Gly Ile Pro Glu Pro Cys  
 40  
 384 409 424 439  
 TGT GTG CCA GAA AAG AAT TCC TCA CTC AAG ATC TTA TTC TTT GAT GAA AAG AAG  
 Cys Val Pro Glu Lys Met Ser Ser Leu Ser Ile Leu Phe Phe Asp Glu Asn Lys  
 45  
 454 459 464  
 AAT GTC GGA CTT AAA GGA TAT CCA AAC AAG AAG GGA GAG TGT TGT GAT TCC AAG  
 Asn Val Val Leu Lys Val Tyr Pro Asn Met Thr Val Glu Ser Cys Ala Cys Arg  
 50  
 503 513 523 533  
 TACCTGAG AAGGAGAT CCGAGCTG AATGAGAG

2. Gène codant pour la BMP-3 bovine ayant la séquence d'acides aminés donnée à la revendication 1.

3. Gène codant pour une protéine montrant au moins les propriétés de la BMP-3 pour induire la formation osseuse et comprenant une séquence d'ADN :

(a) qui diffère d'une séquence d'ADN de la revendication 1 dans la séquence de codons du fait de la dégénérescence du code génétique ;



(b) qui s'hybride avec une séquence d'ADN de la revendication 1 ou du paragraphe (a) ci-dessus dans des conditions d'hybridation stringentes; ou  
 (c) représente un fragment ou une variation allélique d'une séquence d'ADN de la revendication 1.

- 5 4. Séquence d'ADN suivant la revendication 3, qui code pour la BMP-3 humaine.
5. Séquence d'ADN suivant la revendication 3 ou 4, qui est une séquence d'ADN génomique.
6. Séquence d'ADN suivant la revendication 3 ou 4, qui est une séquence d'ADNc.
- 10 7. Vecteur contenant le gène ou la séquence d'ADN suivant l'une quelconque des revendications 1 à 6, en association opérationnelle avec une séquence de contrôle d'expression.
8. Cellule transformée avec un vecteur de la revendication 7.
- 15 9. Cellule suivant la revendication 8, qui est une cellule mammifère, une cellule bactérienne, une cellule d'insecte ou une cellule de levure.
10. Cellule suivant la revendication 9, qui est une cellule CHO.
- 20 11. Protéine montrant des propriétés de la BMP-3, qui est codée par le gène ou la séquence d'ADN de l'une quelconque des revendications 1 à 6.
- 25 12. Protéine montrant des propriétés de la BMP-3, qui est produite par les étapes de culture dans un milieu de culture approprié d'une cellule transformée avec un vecteur d'expression comprenant un gène ou une séquence d'ADN de l'une quelconque des revendications 1 à 6, et de récupération de ladite protéine du milieu de culture précité.
- 30 13. Procédé de production de la protéine suivant l'une ou l'autre des revendications 11 ou 12, comprenant les étapes de culture dans un milieu de culture approprié de la cellule de la revendication 9 et d'isolement de ladite protéine du milieu de culture précité.
14. Composition pharmaceutique comprenant la protéine suivant l'une ou l'autre des revendications 11 ou 12 et un véhicule pharmaceutiquement acceptable.
- 35 15. Composition pharmaceutique suivant la revendication 14, comprenant en outre une matrice pouvant distribuer la composition au site de l'anomalie osseuse ou cartilagineuse et fournir une structure pour induire une formation osseuse ou cartilagineuse.
- 40 16. Composition pharmaceutique suivant la revendication 15, dans laquelle ladite matrice comprend de l'hydroxyapatite, du collagène, de l'acide polylactique ou du phosphate tricalcique.
17. Utilisation d'une protéine suivant l'une ou l'autre des revendications 11 ou 12 pour la préparation d'une composition pharmaceutique pour induire une formation osseuse ou cartilagineuse.

#### Revendications pour l'Etat contractant suivant : AT

1. Procédé de préparation d'un gène codant pour le BMP-3 bovin comprenant la séquence d'ADN suivante :

383 393 403 413 423  
 GAGGAGAG GAGGAGAG GAGGAGAG GAGGAGAG AAC AAT GAG CTT CTT GGG GGA  
 Asn Asn Glu Leu Pro Gly Ala  
 5  
 443 453 473 483  
 GAA TTT CAG TAC AAG GAG GAT GAA GAA TGG GAG GAG AAG AAG CTT TAC AAG ACT  
 Glu Tyr Gln Tyr Lys Glu Asp Glu Val Trp Glu Glu Arg Lys Pro Tyr Lys Thr  
 10  
 503 513 533  
 CTT GAG ACT CAG CCC CTT GAT AAG AAT AAG AAC AAA AAG AAA CAG AAG AAG GGA  
 Leu Gln Thr Gln Pro Pro Asp Lys Ser Lys Asn Lys Lys Lys Gln Arg Lys Gly  
 15  
 543 563 573 583  
 CTT GAG CAG AAG AAT CAG AAG CTC CAG TTT GAT GAA CAG AAT CTC AAG AAG GGA  
 Pro Gln Gln Lys Ser Gln Thr Leu Gln Phe Asp Glu Gln Thr Leu Lys Lys Ala  
 20  
 603 623 633  
 AAG AAG AAG CAA TGG AAT GAA CCC CAG AAT TGT GGC AAG CAG TAC CTT AAA GTG  
 Arg Arg Lys Gln Trp Ile Glu Pro Arg Asn Cys Ala Arg Arg Tyr Leu Lys Val  
 25  
 653 663 683 693  
 GAC TTC GGA GAT AAT GGC TGG AAG GAA TGG AAT AAT TCC CCC AAG TCC TTC GAT  
 Asp Phe Ala Asp Ile Gly Trp Ser Glu Trp Ile Ile Ser Pro Lys Ser Phe Asp  
 713 723 743 753 763  
 GGC TAT TAC TGC TCC GGA GGC TCC CAG TTC CAC AAG CCA AAG GAGGAGAG TTTTGGCC--  
 Ala Tyr Tyr Cys Ser Gly Ala Cys Gln Phe Pro Met Pro Lys  
 773 783  
 TGTCTTCCG ATTTCCTAG : et  
 30  
 284 294 304 313  
 CTAAGTGGG TTTCTCTTT TGTCTCTAG TCT TTC AAG CCA TCA AAT CAG GGT ACC  
 Ser Leu Lys Pro Ser Asn His Ala Thr  
 35  
 334 349 354 373  
 ATC CAG AAT AAT GAG AAG GGT GTG GGG GTC GTC CTT GGA ATC CTT GAG CTT TGC  
 Ile Gln Ser Ile Val Arg Ala Val Gly Val Val Pro Gly Ile Pro Glu Pro Cys  
 40  
 384 409 424 439  
 TGT GTC CCA GAA AAG ATG TCC TCA CTC AAG ATC TTA TTC TTT GAT GAA AAG AAG  
 Cys Val Pro Glu Lys Met Ser Ser Leu Ser Ile Leu Phe Phe Asp Glu Asn Lys  
 45  
 454 469 484  
 AAT GTC GAA CTT AAA GAA TAT CCA AAC AAG AAG GAA GAG TCT TGT GGT TGC AAG  
 Asn Val Val Leu Lys Val Tyr Pro Asn Met Thr Val Glu Ser Cys Ala Cys Arg  
 503 513 523 533  
 TACCTGGG AAGACTCT CTGCTGCTT AATCTATC.

ledit procédé comprenant le criblage d'une banque génomique recombinante de bovin avec des sondes  
 comprenant des pools d'oligonucléotides conçues sur la base des séquences d'acides aminés partielles et l'iso-  
 lement d'un clone positif.

2. Procédé selon la revendication 1, dans lequel le gène codant pour le BMP-3 a la séquence d'acides aminés donnée dans la revendication 1.
3. Procédé selon la revendication 1, dans lequel un gène codant pour une protéine présentant au moins la propriété

du BMP-3 d'induction de la formation de l'os comprend une séquence d'ADN :

- a) qui diffère d'une séquence d'ADN obtenue selon la procédé de la revendication 1 par la séquence de codons du fait de la dégénérescence du code génétique ;
- b) qui s'hybride avec une séquence d'ADN obtenue selon le procédé de la revendication 1 ou la section a) ci-dessus dans des conditions d'hybridation stringentes ; ou
- c) représente un fragment ou une variation allélique d'une séquence d'ADN obtenue selon le procédé de la revendication 1,

4. Procédé selon la revendication 3, dans lequel la séquence d'ADN code pour le BMP-3 humain.

5. Procédé selon la revendication 3 ou 4, dans lequel la séquence d'ADN est une séquence d'ADN génomique.

6. Procédé selon la revendication 3 ou 4, dans lequel la séquence d'ADN est une séquence d'ADNc.

7. Procédé de préparation d'un vecteur recombinant comprenant l'insertion du gène ou de la séquence d'ADN obtenue selon un procédé de l'une quelconque des revendications 1 à 6 en association opératoire avec une séquence de contrôle d'expression dans un vecteur adapté.

8. Cellule transformés avec un vecteur préparé selon le procédé de la revendication 7.

9. Cellule selon la revendication 8, qui est une cellule mammifère, une cellule bactérienne, une cellule d'insecte ou une cellule de levure.

10. Cellule selon la revendication 9, qui est une cellule CHO.

11. Procédé de production d'une protéine présentant les propriétés du BMP-3, comprenant les étapes de culture dans un milieu de culture adapté de la cellule de la revendication 9 et d'isolement de ladite protéine dudit milieu de culture.

12. Procédé de préparation d'une composition pharmaceutique comprenant la combinaison de la protéine produite selon le procédé de la revendication 11 avec un véhicule pharmaceutiquement acceptable.

13. Procédé selon la revendication 12, dans lequel la composition pharmaceutique comprend en outre une matrice capable de délivrer la composition sur le site du défaut de l'os ou du cartilage et la fourniture d'une structure pour induire la formation d'os ou de cartilage.

14. Procédé selon la revendication 13, dans lequel ladite matrice de ladite composition pharmaceutique comprend de l'hydroxyapatite, du collagène, de l'acide polylactique ou du phosphate tricalcique.

15. Utilisation d'une protéine selon la revendication 11 pour la préparation d'une composition pharmaceutique pour induire la formation d'os ou de cartilage.

16. Gène codant pour le BMP-3 bovin comprenant la séquence d'ADN suivante :

55      **17. Gène codant pour le BMP-3 bovin ayant la séquence d'acides aminés donnée dans la revendication 16.**

**18. Gène codant pour une protéine présentant au moins la propriété du BMP-3 d'induction de la formation de l'es et comprenant une séquence d'ADN :**



- a) qui diffère d'une séquence d'ADN de la revendication 16 par la séquence de codons du fait de la dégénérescence du code génétique ;  
 b) qui s'hybride avec une séquence d'ADN de la revendication 16 ou de la section a) ci-dessus dans des conditions d'hybridation stringentes ou  
 c) représente un fragment ou une variation allélique d'une séquence d'ADN selon la revendication 16.

19. Séquence d'ADN selon la revendication 18, qui code pour le BMP-3 humain.

20. Séquence d'ADN selon la revendication 18 ou 19, qui est une séquence d'ADN génomique.

21. Séquence d'ADN selon la revendication 18 ou 19, qui est une séquence d'ADNc.

22. Vecteur contenant le gène ou la séquence d'ADN selon l'une quelconque des revendications 16 à 21 en association opératoire avec une séquence de contrôle d'expression.

23. Cellule transformée avec un vecteur selon la revendication 22.

24. Cellule selon la revendication 23, qui est une cellule mammifère, une cellule bactérienne, une cellule d'insecte ou une cellule de levure,

25. Cellule selon la revendication 24, qui est une cellule CHC.

26. Protéine présentant les propriétés du BMP-3 qui est encodée par le gène ou la séquence d'ADN selon l'une quelconque des revendications 1 à 21.

27. protéine présentant les propriétés du BMP-3 qui est produite par les étapes de culture dans un milieu de culture adapté d'une cellule transformée avec un vecteur d'expression comprenant un gène ou la séquence d'ADN selon l'une quelconque des revendications 1 à 21, et de récupération de ladite protéine dudit milieu de culture.

28. Composition pharmaceutique comprenant la protéine selon la revendication 26 ou 27 et un véhicule pharmaceutiquement acceptable.

29. Composition pharmaceutique selon la revendication 29, comprenant en outre une matrice capable de délivrer la composition sur le site du défaut de l'os ou du cartilage et la fourniture d'une structure pour induire la formation d'os ou de cartilage.

30. Composition pharmaceutique selon la revendication 29, dans laquelle ladite matrice comprend de l'hydroxyapatite, du collagène, de l'acide polylactique ou du phosphate tricalcique.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**